

UDK 517.929.2:517.929.4

SCHEME OF THE CHAIN METHOD

Kruglov Viktor

*c. ph.-m. s., prof., professor of the Department
of Methods of Mathematical Physics
Odessa I.I. Mechnikov National University,
Odessa, str. Dvoryanska, 2, 65029
Odessa, Ukraine*

Анотація. В цій роботі наведено схему методу ланцюгів стосовно розв'язання скінченного лінійного різницевого рівняння, і приведено формулу загального розв'язку цього рівняння. Як наслідок, наведено формулу загального розв'язку різницевого рівняння зі сталими коефіцієнтами, яка цілком залежить тільки від коефіцієнтів цього рівняння. Розглянуті розв'язки лінійних диференціальних рівнянь у вигляді узагальненого степеневого ряду, коефіцієнти якого знаходяться методом ланцюгів.

Ключові слова: ланцюг, різницеве рівняння, функція цілочисельного аргументу.

Abstract. A scheme of the chain method for solving a finite linear difference equation given in this paper, and a formula for this equation's general solution of is given. As a result, the formula for the general solution of a difference equation with constant coefficients is given. This formula depend entirely only on the coefficients of this equation. Considered solutions of linear differential equations in the form of a generalized power series, the coefficients of which are found by the chain method.

Keywords: chain, difference equation, functions of integer argument.

Introduction.

This paper is a review of the author's works [1 – 5], in which the chain method was developed for solving a linear difference equation and applied to construct solutions to some linear differential equations.

Main text.

Consider a linear difference equation of order n :

$$l_{n+k} = a_{1k}l_{n+k-1} + a_{2k}l_{n+k-2} + \dots + a_{nk}l_k, \quad a_{nk} \neq 0, \quad k = 0, 1, \dots, \quad (1)$$

where $a_{1k}, a_{2k}, \dots, a_{nk}$ are known functions of integer argument k .

A step-by-step solution of equation (1) is used, i.e. at each subsequent step of using equation (1), the solutions found in the previous steps are taken into account, and writing the coefficients of equation (1) in the form a_{jk} does not allow establishing the regularities that exist between these coefficients. Therefore, we denote $a_{jk} = a_{n+k-j}^{(j)}$, $j = \overline{1, n}$. Then equation (1) takes the form

$$l_{n+k} = a_{n+k-1}^{(1)}l_{n+k-1} + a_{n+k-2}^{(2)}l_{n+k-2} + \dots + a_k^{(n)}l_k, \quad k = 0, 1, \dots. \quad (2)$$

A chain consisting of elements of the set $M_{n,n+k}$ is called the product of the maximum possible number of elements from this set, but for two arbitrary adjacent factors in this product the specified multiplication rule must be used. It follows that an arbitrary chain starts with some initial element of the set $M_{n,n+k}$ and ends with one of the final elements of this set.

The structure of an arbitrary chain has the following form:

$$a_n^{(i_1)} a_{n+i_1}^{(i_2)} a_{n+i_1+i_2}^{(i_3)} \cdot \dots \cdot a_{n+i_1+i_2+\dots+i_{r-1}}^{(i_r)},$$

$$n + i_1 + \dots + i_{r-1} + i_r = n + k.$$

The order of a chain is the sum of the ranks of all the factors that form this chain. From the elements of the set $M_{n,n+k}$ it is possible to form chains of order k only, because $i_1 + \dots + i_{r-1} + i_r = k$.

The function $f_{k,n+k}$ is the sum of all chains of order k that can be composed of elements of the set $M_{n,n+k}$. Let us now count the number of terms in the function $f_{k,n+k}$. Let a chain of order k have x_1 elements of rank one, x_2 elements of rank two, etc., x_n elements of rank n . Then $k = x_1 + 2x_2 + \dots + nx_n$. It follows that the number of all chains of order k that can be composed of elements of the set $M_{n,n+k}$ is equal to

$$Q_k^{(n)} = \sum_{x_1+2x_2+\dots+nx_n=k} \frac{(x_1 + x_2 + \dots + x_n)!}{x_1! x_2! \dots x_n!}.$$

The function $f_{k-1,n+k}$ is constructed similarly. It is the sum of chains of order $k - 1$, which are formed from the elements of the set

$$M_{n+1,n+k} = \left(a_{n+1}^{(1)}, \dots, a_{n+k-1}^{(1)}; a_{n+1}^{(2)}, \dots, a_{n+k-2}^{(2)}; \dots; a_{n+1}^{(p)}, \dots, a_q^{(p)} \right),$$

and the number of such chains is

$$Q_{k-1}^{(n)} = \sum_{x_1+2x_2+\dots+nx_n=k-1} \frac{(x_1+x_2+\dots+x_n)!}{x_1!x_2!\dots x_n!}, \text{ etc.}$$

The number of chains of order $k - m$ (i.e., terms in the function $f_{k-m,n+k}$) formed from the elements of the set

$$M_{n+m,n+k} = \left(a_{n+m}^{(1)}, \dots, a_{n+k-1}^{(1)}; a_{n+m}^{(2)}, \dots, a_{n+k-2}^{(2)}; \dots; a_{n+m}^{(p)}, \dots, a_q^{(p)} \right),$$

is equal to

$$Q_{k-m}^{(n)} = \sum_{x_1+2x_2+\dots+nx_n=k-m} \frac{(x_1 + x_2 + \dots + x_n)!}{x_1! x_2! \dots x_n!}.$$

The solution to equation (2) is given by formulas (3) and (4).

Let

$$a_{jk} = a_{n+k-j}^{(j)} \equiv a_j, \quad k = 0, 1, \dots,$$

that is, we are dealing with a difference equation with constant coefficients

$$l_{n+k} = a_1 l_{n+k-1} + a_2 l_{n+k-2} + \dots + a_n l_k, \quad k = 0, 1, \dots \quad (6)$$

Now, in any chain, the order of multiplication of its elements according to the specified rule loses its meaning, and the chain structure takes the form $a_1^{x_1} a_2^{x_2} \dots a_n^{x_n}$.

Then

$$f_{k,n+k} = \sum_{x_1+2x_2+\dots+nx_n=k} \frac{(x_1 + x_2 + \dots + x_n)!}{x_1! x_2! \dots x_n!} \cdot a_1^{x_1} a_2^{x_2} \dots a_n^{x_n} = R_k^{(n)},$$

$$f_{k-m,n+k} = \sum_{x_1+2x_2+\dots+nx_n=k-m} \frac{(x_1 + x_2 + \dots + x_n)!}{x_1! x_2! \dots x_n!} \cdot a_1^{x_1} a_2^{x_2} \dots a_n^{x_n} = R_{k-m}^{(n)}.$$

Equation (5) can be written in the following way:

$$R_k^{(n)} = a_1 R_{k-1}^{(n)} + a_2 R_{k-2}^{(n)} + \dots + a_n R_{k-n}^{(n)}, \quad k > n;$$

$$R_k^{(n)} = a_1 R_{k-1}^{(n)} + \dots + a_{k-1} R_1^{(n)}, \quad k = \overline{1, n}; \quad (7)$$

$$R_0^{(n)} = 1, R_1^{(n)} = a_1. \quad (8)$$

Equation (4) can be written in the following way:

$$\varphi_{0,n+k} = a_n R_k^{(n)},$$

.....

$$\varphi_{i,n+k} = a_{n-i} R_k^{(n)} + \dots + a_n R_{k-i}^{(n)}, \quad i = \overline{1, n-2},$$

.....

$$\varphi_{n-1,n+k} = R_{k+1}^{(n)} = a_1 R_k^{(n)} + \dots + a_n R_{k-n+1}^{(n)}.$$

The general solution of equation (6) is given by the formula

$$l_{n+k} = \sum_{i=1}^n (a_i R_k^{(n)} + a_{i+1} R_{k-1}^{(n)} + \dots + a_n R_{k-n+i}^{(n)}) l_{n-i}, \quad k = 0, 1, \dots, \quad (9)$$

where the numbers $R_k^{(n)}$, if $k = \overline{1, n}$ or $k > n$, are calculated by formulas (7), (8).

It can be proven that among the solutions (9) there are known solutions of the

form $k^m \lambda^s$, where λ is the root of the characteristic equation

$$\lambda^n = a_1 \lambda^{n-1} + \dots + a_{n-1} \lambda + a_n.$$

In [1] an example is given where the equation is considered $l_{k+2} = al_{k+1} + bl_k$.

Its solution is given by the formula

$$l_{k+2} = (al_1 + bl_0) \sum_{m=0}^{[k/2]} C_{k-m}^m a^{k-2m} b^m + bl_1 \sum_{m=0}^{[k/2]} C_{k-m-1}^m a^{k-2m-1} b^m.$$

This formula implies:

- 1) if $\lambda_1 \neq \lambda_2$ and $l_1 = \lambda_1, l_0 = 1$, then $l_{k+2} = \lambda_1^{k+2}$;
- 2) if $\lambda_1 = \lambda_2 = \frac{a}{2}$, $l_1 = \lambda_1 = \frac{a}{2}, l_0 = 1$, then

$$l_{k+2} = \left(\frac{a}{2}\right)^{k+2} \lambda_1^{k+2},$$

and for $l_1 = \lambda_1, l_0 = 0$

$$l_{k+2} = (k+2) \lambda_1^{k+2}.$$

Summary and conclusions.

Thus, the construction of a fundamental system for solving equation (2) allows us to explicitly construct a general solution to a homogeneous difference equation of the n -th order. In this paper, we consider the case where the coefficients of the difference equation are constant real numbers. The second-order equations are generalized by the Fibonacci sequence.

References:

1. Kruglov V.E. (2009). Construction of a fundamental system of solution of a linear finite-order difference equation. *UMJ*, Vol. 61(6), Pp. 923 – 944.
2. Kruglov V.E. (2016). On n -arithmetical triangles constructed for polynomial coefficients. *RM*, Vol. 60(8), P. 29.
3. Kruglov V.E. (2008). Solution of a second-order Poincare-Perron-type equation and differential equations that can be reduced to it. *UMJ*, Vol. 60(7), Pp. 1055 – 1071.
4. Kruglov V.E. (2010). Solution of the linear differential equation of n -th order

with four singular points. *Annales Univ. Sci. Budapest, Sect. Comp.*, Vol. 32, Pp. 23 – 35.

5. Kruglov V.E. (2013). Solution of the linear second-order differential equation with coefficients analytic in the vicinities of a Fuchsian zero point. *UMJ*, Vol (64(10)), Pp. 1572 – 1585.

6. Kruglov V.E. (2023). The chain method for solving a finite-order linear difference equation and some of its applications. *Researches in Mathematics and Mechanics*. Vol. 28. Issue 1 – 2 (41 – 42), Pp. 40 – 46.

The article has been sent: 25.11.2025.

© Kruglov V.E.